

Novel Electrochemistry of The Ru(0001) Surface Towards Electrocatalysis At Molecular Level

W. F. Lin, P. A. Christensen and A. Hamnett

Department of Chemistry, Bedson Building, The University, Newcastle upon Tyne, NE1 7RU, UK. Tel: +44-191-2227056; Fax: +44-191-2226929; Email: wenfeng.lin@ncl.ac.uk

which oxidation of CO_L was observed, suggesting a higher mobility of the CO_L species formed from the adsorption of the HCOOH . At potentials > 1000 mV, the oxidation of formic acid to CO_2 was significantly increased, and the oxidation of methanol to CO_2 and methyl formate was observed, both of which were attributed to the formation of an active RuO_2 phase on the Ru(0001) surface.

Abstract

Ex-situ LEED, RHEED and Auger data have shown that the Ru(0001) electrode shows the potential-dependent formation of well-defined and ordered oxygen-containing adlayers of (2×2) , (3×1) and $(1 \times 1)\text{-O}$ and finally $\text{RuO}_2(100)$ islands in perchloric acid solution. The adsorption and electro-oxidation of methanol, formic acid and CO at the Ru(0001) electrode have been investigated as a function of temperature, potential and time using *in-situ* FTIR spectroscopy, and the results interpreted in terms of the surface chemistry of the Ru(0001) electrode. It was found that the oxide phases have a marked effect upon the reactivity of the surface towards the electro-oxidation of such fuel molecules; in addition, it is clear from the data that RuO_2 is significantly active towards the oxidation of both methanol and formic acid.

It was found that no dissociative adsorption or electro-oxidation of methanol takes place at the Ru(0001) at potentials < 800 mV vs Ag/AgCl, and at all three temperatures employed, 10°C , 25°C and 50°C . However, formic acid did undergo dissociative adsorption, even at -200 mV, to form linearly adsorbed CO, (CO_L), with little or no CO adsorbed at threefold-hollow sites, (CO_H). In contrast to the adsorption of CO, it was found that increasing the temperature to 50°C markedly increased the amount of CO_L formed on the Ru(0001) surface from the adsorption of formic acid. On increasing the potential, the electro-oxidation of the CO_L species to CO_2 took place *via* reaction with the active $(3 \times 1)\text{-O}$ oxide. At all three temperatures employed, it was found that the formic acid was oxidised to CO_2 faster than was observed in the experiments involving CO adsorbed from $\text{CO}(\text{g})$ over the potential range at